

Monoferric[™] (ferric derisomaltose injection)

(Intravenous)

Document Number: EOCCO-0524

Last Review Date: 05/02/2024 Date of Origin: 02/04/2020 Dates Reviewed: 02/2020, 07/2020, 10/2020, 12/2021, 12/2022, 12/2023, 05/2024

I. Length of Authorization

Coverage will be provided for 35 days.

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC unit]:

- Monoferric 100 mg/1 mL single-dose vial: 4 vials per 35 days
- Monoferric 500 mg/5 mL single-dose vial: 1 vial per 35 days
- Monoferric 1000 mg/10 mL single-dose vial: 1 vial per 35 days

B. Max Units (per dose and over time) [HCPCS Unit]:

• 100 billable units per 35 days

III. Initial Approval Criteria ¹⁻¹¹

Coverage is provided in the following conditions:

- Patient had an inadequate response, or has a contraindication or intolerance, to sodium ferric gluconate complex (Ferrlecit[®]) OR iron dextran (INFeD[®]) OR iron sucrose (Venofer[®]); **AND**
- Patient is at least 18 years of age; AND
- Laboratory values must be obtained within 28 days prior to the anticipated date of administration; **AND**
- Other causes of anemia (e.g., vitamin B-12 deficiency, thalassemia, sideroblastic anemia, etc.) have been ruled out; **AND**
- Patient does not have a history of allergic reaction to any intravenous iron product; AND
- Other supplemental iron is to be discontinued prior to administration of ferric derisomaltose; **AND**

Iron Deficiency Anemia in Non-Dialysis-Dependent Chronic Kidney Disease (NDD-CKD) † ^{1,10,11}

• Patient must not be receiving hemodialysis; AND



- Patient has chronic renal impairment with eGFR between 15-59 mL/min; AND
- Patient has iron-deficiency anemia with a Hemoglobin (Hb) ≤11 g/dL; AND
 - Ferritin ≤100 ng/mL; **OR**
 - Ferritin \leq 300 ng/mL when transferrin saturation (TSAT) \leq 30%

Iron Deficiency Anemia in patients intolerant to or who have had unsatisfactory response to oral iron † ^{1,9}

- Patient had an intolerance or inadequate response to a minimum of 14 days of oral iron; AND
- Patient has iron-deficiency anemia with a Hemoglobin (Hb) ≤11 g/dL; AND
 - Ferritin <100 ng/mL; AND
 - Transferrin saturation (TSAT) <20%

Cancer- and Chemotherapy-Induced Anemia ‡ ^{5,6}

- Used as a single agent; AND
 - Patient has absolute iron deficiency defined as ferritin < 30 ng/mL <u>AND</u> a TSAT < 20%;
 OR
 - Patient has functional iron deficiency defined as ferritin > 500 800 ng/mL <u>AND</u> a TSAT
 < 50% with the goal of avoiding allogenic transfusion; **OR**
- Used in combination with erythropoiesis-stimulating agents (ESAs); AND
 - Patient has absolute iron deficiency defined as ferritin < 30 ng/mL <u>AND</u> a TSAT < 20% and failed to demonstrate an increase in Hb after 4 weeks of IV or oral iron therapy; **OR**
 - Patient has functional iron deficiency defined as ferritin 30 500 ng/mL <u>AND</u> a TSAT < 50% and is receiving myelosuppressive chemotherapy without curative intent

† FDA Approved Indication(s); **‡** Compendia Recommended Indication(s); **Φ** Orphan Drug

IV. Renewal Criteria ^{1,5}

Refer to initiation criteria.

V. Dosage/Administration ^{1,5}

Indication	Dose
All indications	Weight \geq 50 kg:
	• Administer 1,000 mg intravenously as a single dose.
	Weight < 50 kg:
	• Administer 20 mg/kg actual body weight intravenously as a single dose.
	Note: Treatment may be repeated if iron deficiency anemia reoccurs.



VI. Billing Code/Availability Information

HCPCS Code:

• J1437 – Injection, ferric derisomaltose, 10 mg: 1 billable unit = 10 mg

NDC(s):

- Monoferric 100 mg/1 mL single-dose vial: 73594-9301-xx
- Monoferric 500 mg/5 mL single-dose vial: 73594-9305-xx
- Monoferric 1000 mg/10 mL single-dose vial: 73594-9310-xx

VII. References

- 1. Monoferric [package insert]. Holbaek, Denmark; Pharmacosmos, A/S. August 2022. Accessed April 2024.
- KDOQI; National Kidney Foundation. Clinical practice guidelines and clinical practice recommendations for anemia in chronic kidney disease in adults. Am J Kidney Dis. 2006 May;47(5 Suppl 3):S16-85.
- Kidney Disease: Improving Global Outcomes (KDIGO) Anemia Work Group. KDIGO Clinical Practice Guideline for Anemia in Chronic Kidney Disease. Kidney inter., Suppl. 2012; 2: 279–335.
- Ratcliffe LE, Thomas W, Glen J, et al. Diagnosis and Management of Iron Deficiency in CKD: A Summary of the NICE Guideline Recommendations and Their Rationale. Am J Kidney Dis. 2016 Apr;67(4):548-58.
- 5. Referenced with permission from the NCCN Drugs and Biologics Compendium (NCCN Compendium[®]) ferric derisomaltose. National Comprehensive Cancer Network, 2024. The NCCN Compendium[®] is a derivative work of the NCCN Guidelines[®]. NATIONAL COMPREHENSIVE CANCER NETWORK[®], NCCN[®], and NCCN GUIDELINES[®] are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2024.
- 6. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines[®]) Hematopoietic Growth Factors 3.2024. National Comprehensive Cancer Network, 2024. The NCCN Compendium[®] is a derivative work of the NCCN Guidelines[®]. NATIONAL COMPREHENSIVE CANCER NETWORK[®], NCCN[®], and NCCN GUIDELINES[®] are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2024.
- Wish JB. Assessing iron status: beyond serum ferritin and transferrin saturation. Clin J Am Soc Nephrol. 2006 Sep;1 Suppl 1:S4-8.
- 8. Koch TA, Myers J, Goodnough LT. Intravenous Iron Therapy in Patients with Iron Deficiency Anemia: Dosing Considerations. Anemia. 2015;2015:763576.



- Auerbach M, Henry D, Derman RJ, et al. A prospective, multi-center, randomized comparison of iron isomaltoside 1000 versus iron sucrose in patients with iron deficiency anemia; the FERWON-IDA trial. Amer J of Hema. Sep2019:94;9;pps1007-1014.
- Bhandari S, Thomsen LL. Single 1000 Mg Infusion Of Iron Isomaltoside 1000 Demonstrates A More Rapid Hemoglobin Response And Reduced Risk Of Cardio-Vascular Adverse Events Compared To Multiple Doses Of IV Iron Sucrose In The FERWON Trials. Nephrology Dialysis Transplantation 34 (Supplement 1): i475–i486, 2019.
- Bhandari S, Kalra PA, Berkowitz M, et al. Safety and efficacy of iron isomaltoside 1000/ferric derisomaltose versus iron sucrose in patients with chronic kidney disease: the FERWON-NEPHRO randomized, open-label, comparative trial. Nephrol Dial Transplant. 2021 Jan 1;36(1):111-120. doi: 10.1093/ndt/gfaa011.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
D50.0	Iron deficiency anemia secondary to blood loss (chronic)
D50.1	Sideropenic dysphagia
D50.8	Other iron deficiency anemias
D50.9	Iron deficiency anemia, unspecified
D63.0	Anemia in neoplastic disease
D63.1	Anemia in chronic kidney disease
D63.8	Anemia in other chronic disease classified elsewhere
D64.81	Anemia due to antineoplastic chemotherapy
Z51.11	Encounter for antineoplastic chemotherapy
Z51.89	Encounter for other specified aftercare

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

The preceding information is intended for non-Medicare coverage determinations. Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determinations (NCDs) and/or Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. Local Coverage Articles (LCAs) may also exist for claims payment purposes or to clarify benefit eligibility under Part B for drugs which may be self-administered. The following link may be used to search for NCD, LCD, or LCA documents: https://www.cms.gov/medicare-coverage-database/search.aspx. Additional indications, including any preceding information, may be applied at the discretion of the health plan.



Medicare Part B Administrative Contractor (MAC) Jurisdictions			
Jurisdiction	Applicable State/US Territory	Contractor	
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC	
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC	
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)	
6	MN, WI, IL	National Government Services, Inc. (NGS)	
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.	
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)	
N (9)	FL, PR, VI	First Coast Service Options, Inc.	
J (10)	TN, GA, AL	Palmetto GBA	
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA	
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.	
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)	
15	кү, он	CGS Administrators, LLC	

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCA/LCD): N/A