

## Substance Use Disorder Diagnosis Coding Guide for Initiation & Engagement of Substance Use Disorder Treatment (IET) Measure

The IET denominator includes patients with diagnoses of substance <u>abuse</u> or <u>dependence</u>. Patients with diagnoses of SUD <u>recovery</u>, <u>remission</u>, or <u>use</u> do not count in the denominator. It is important for providers, billers, and clinic staff to use accurate diagnosis codes to ensure that patients who do *not* abuse or depend on substances are *not* included in the measure denominator. Following this guide will also ensure that patients who s hould be in the measure denominator are included and that any subsequent SUD treatment services are counted toward the numerator(s).

I	If a	natient is	in long-	term re	covery o	r remission	from SUD
ı	II U	pulicitis	III IOIIG		covciv	1 1 61111331011	110111300

Instead of using these codes, which count toward the IET denominator:		Consider using:	
F10.10	Alcohol abuse, uncomplicated	F10.11	Alcohol abuse, in remission
F10.20	Alcohol dependence, uncomplicated	F10.21	Alcohol dependence, in remission
F11.10	Opioid abuse, uncomplicated	F11.11	Opioid abuse, in remission
F11.20	Opioid dependence, uncomplicated	F11.21	Opioid dependence, in remission
F12.10	Cannabis abuse, uncomplicated	F12.11	Cannabis abuse, in remission
F12.20	Cannabis dependence, uncomplicated	F12.21	Cannabis dependence, in remission
F15.10	Other stimulant abuse, uncomplicated	F15.11	Other stimulant abuse, in remission
F15.20	Other stimulant dependence, uncomplicated	F15.21	Other stimulant dependence, in remission
F19.10	Other psychoactive substance abuse, uncomplicated	F19.11	Other psychoactive substance abuse, in remission
F19.20	Other psychoactive substance dependence, uncomplicated	F19.21	Other psychoactive substance dependence, in remission

## If a patient uses a substance but does not demonstrate abuse or dependence...

Instead of u	sing these codes, which count toward the IET denominator:	Consider using:		
F10.10	Alcohol abuse, uncomplicated	F10.90	Alcohol use, unspecified, uncomplicated	
F10.20	Alcohol dependence, uncomplicated		Alconor use, unspecified, uncomplicated	
F11.10	Opioid abuse, uncomplicated	F11.90	Opioid use, unspecified, uncomplicated	
F11.20	Opioid dependence, uncomplicated	111.90	Opiola use, unspecified, uncomplicated	
F12.10	Cannabis abuse, uncomplicated	F12.90	Cannabis use, unspecified, uncomplicated	
F12.20	Cannabis dependence, uncomplicated	112.50		
F15.10	Other stimulant abuse, uncomplicated		Other stimulant use, unspecified, uncomplicated	
F15.20	Other stimulant dependence, uncomplicated	F15.90	Other stimulant use, unspecified, uncomplicated	
F19.10	Other psychoactive substance abuse, uncomplicated	F19.90	Other psychoactive substance use, unspecified, uncomplicated	
F19.20	Other psychoactive substance dependence, uncomplicated		other psychoactive substance use, unspecified, uncomplicated	

## Please Note:

- > The document is meant for educational purposes only. Clinicians and billers should always use their best judgment on which diagnosis code(s) are clinically appropriate.
- > This is not an exhaustive list. Please reach out to EOCCOmetrics@modahealth.com for a full list of qualifying diagnosis codes.